

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



(2F) ✓

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 48/00	A1	(11) International Publication Number: WO 93/09815 (43) International Publication Date: 27 May 1993 (27.05.93)
(21) International Application Number: PCT/US92/09849 (22) International Filing Date: 19 November 1992 (19.11.92) (30) Priority data: 797,555 22 November 1991 (22.11.91) US (71) Applicants: THE GENERAL HOSPITAL CORPORATION [US/US]; 55 Fruit Street, Boston, MA 02114 (US). THE UNITED STATES OF AMERICA represented by THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES [US/US]; c/o National Institute of Health, Office of Technology Transfer, Box OTT, Bethesda, MD 20892-9902 (US). (72) Inventor: SACHS, David, H. ; 77 Studio Road, Newton, MA 02166 (US).		(74) Agent: CLARK, Paul, T.; Fish & Richardson, 225 Franklin Street, Boston, MA 02110-2804 (US). (81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, SE). Published <i>With international search report.</i> AML INFORMATION SERVICES P.O. BOX 405, CORTE MADERA, CA 94976-0405 (415) 927-0340 • FAX (415) 927-7250
(54) Title: SPECIFIC TOLERANCE IN TRANSPLANTATION		
(57) Abstract		
Use of nucleic acid encoding a mammal MHC antigen inserted into a bone marrow hematopoietic stem cell in the preparation of a medicament for conferring tolerance to a transplanted tissue in a recipient mammal.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LJ	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TG	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

- 1 -

SPECIFIC TOLERANCE IN TRANSPLANTATIONBackground of the Invention

The invention relates to conferring tolerance to
5 foreign tissue.

Tolerance to self major histocompatibility (MHC) antigens occurs during T cell maturation in the thymus (McDuffie et al., 1988, *J. Immunol.* 141:1840). It has been known since the landmark experiments of Billingham,
10 Brent and Medawar (Billingham et al., 1953, *Nat.* 172:603) that exposure of the immune system to allogeneic MHC antigens during ontogeny can cause the immune system to lose reactivity to those antigens, thus leaving the animal specifically tolerant into adult life. Ever since
15 that time, transplantation immunologists have sought means of inducing tolerance in adult animals by production of lymphohematopoietic chimeras. The induction of tolerance across MHC barriers in adult mice by whole body irradiation (WBI) and bone marrow
20 transplantation (BMT) has been studied extensively in murine models (Rayfield et al., 1983, *Transplan.* 36:183; Mayumi et al., 1989, *J. Exp. Med.* 169:213; Sykes et al., 1988, *Immunol. Today* 9:23). This approach has also recently been extended to the miniature swine large
25 animal model, in which it was demonstrated that bone marrow transplants across MHC barriers led to the induction of long-term, specific transplantation tolerance to kidney grafts from donors MHC matched to the bone marrow donors (Guzzetta et al., 1991, *Transplan.*
30 51:862).

The use of MHC mismatched BMT as a means of inducing tolerance to organ allografts is usually accompanied by several major disadvantages: the preparative regimen required for allogeneic BMT involves
35 lethal irradiation, with its inherent risks and

- 2 -

toxicities; clinical applicability is limited by the fact that most potential recipients do not have an appropriate MHC-matched donor, and BMT across MHC barriers causes severe graft-vs-host disease (GVHD); and removing the T lymphocytes in allogeneic bone marrow inocula (Radt et al., 1971, *Eur. J. Immunol.* 4:25), to prevent GVHD is associated with increased rates of engraftment failure (Martin et al., 1988, *Bone Marrow Transplant* 3:445; O'Reilly et al., 1985, *Transplant. Proc.* 17:455; Soderling et al., 1985, *J. Immunol.* 135:941). While these drawbacks are generally considered acceptable for the treatment of otherwise lethal malignant diseases (e.g., leukemia), they would severely limit the application of this methodology as a preparative regimen for organ transplantation, in which non-specific immunosuppressive agents, while not without major complications, are nevertheless effective.

There have been previous reports of the use of cell lines transfected with allogeneic class I and class II genes to selectively modify the immune response to subsequent tissue grafts bearing the foreign gene (Madsen et al., 1989, *Transplant. Proc.* 21:477). Graft prolongation by this technique was not permanent, and the mechanism is unclear. Several laboratories have demonstrated the utility of retroviral mediated gene transfer for the introduction of new genetic material into totipotent hematopoietic stem cells of mice. In general, these protocols involve the transduction of bone marrow by recombinant retroviral vectors ex vivo, with the subsequent reintroduction of the treated cells into myeloablated recipients (for review, see Dick et al., 1986, *Trends in Genetics* 2:165).

- 3 -

Summary of the Invention

In general, the invention features the use of nucleic acid encoding a mammalian MHC antigen inserted into a bone marrow hematopoietic cell in the preparation of a medicament for conferring tolerance to a transplanted tissue in a recipient mammal, e.g., a human.

In preferred embodiments the recipient mammal is from a first species and the transplanted tissue is from a second species; the mammalian MHC antigen is from the second species; the transplanted tissue is from a swine, e.g., a miniature swine; the nucleic acid and the tissue are obtained from the same individual mammal; the nucleic acid is obtained from an individual mammal which is syngeneic to the individual mammal from which the tissue is obtained; the nucleic acid is obtained from an individual mammal which is MHC identical to the individual mammal from which the tissue is obtained; the nucleic acid includes an MHC class I gene; the nucleic acid includes an MHC class II gene; the nucleic acid is inserted into the cell by transduction; the nucleic acid is inserted into the cell by a retrovirus e.g., a Moloney-based retrovirus; the hematopoietic cell is taken from the recipient; the recipient mammal is from a first species and the transplanted tissue is from the first species.

In another aspect, the the invention features the use of a bone marrow hematopoietic cell including a nucleic acid encoding a mammalian MHC antigen in the preparation of a medicament for conferring tolerance to a transplanted tissue in a recipient mammal.

In preferred embodiments the recipient mammal is from a first species and the transplanted tissue is from a second species; the mammalian MHC antigen is from the second species; the transplanted tissue is from a swine, e.g., a miniature swine; the nucleic acid and the tissue

- 4 -

are obtained from the same individual mammal; the nucleic acid is obtained from an individual mammal which is syngeneic to the individual mammal from which the tissue is obtained; the nucleic acid is obtained from an individual mammal which is MHC identical to the individual mammal from which the tissue is obtained; the nucleic acid includes an MHC class I gene; the nucleic acid includes an MHC class II gene; the nucleic acid is inserted into the cell by transduction; the nucleic acid is inserted into the cell by a retrovirus e.g., a Moloney-based retrovirus; the hematopoietic cell is taken from the recipient; the recipient mammal is from a first species and the transplanted tissue is from the first species.

15 In another aspect the invention features a method of inducing tolerance in a recipient mammal, e.g., a human, of a first species to a tissue obtained from a mammal, e.g., a swine, e.g., a miniature swine, of a second species, which tissue expresses an MHC antigen, including inserting DNA encoding an MHC antigen of the second species into a bone marrow hematopoietic stem cell from the recipient mammal, and allowing the MHC antigen encoding DNA to be expressed in the recipient.

Preferred embodiments include those in which: the cell is removed from the recipient mammal prior to the DNA insertion and returned to the recipient mammal after the DNA insertion; the DNA is obtained from the individual mammal from which the tissue is obtained; the DNA is obtained from an individual mammal which is syngeneic with the individual mammal from which the tissue is obtained; the DNA is obtained from an individual mammal which is MHC matched, and preferably identical, with the individual mammal from which the tissue is obtained; the DNA includes an MHC class I gene; the DNA includes an MHC class II gene; the DNA is

- 5 -

insert d into the cell by transduction, e.g., by a retrovirus, e.g., by a Moloney-based retrovirus; and the DNA is expressed in bone marrow cells and/or peripheral blood cells of the recipient 14, preferably 30, more preferably 60, and most preferably 120 days, after the DNA is introduced into the recipient.

In another aspect, the invention features a method of inducing tolerance in a recipient mammal, e.g., a human, to a tissue obtained from a donor mammal of the same species, which tissue expresses an MHC antigen, including: inserting DNA encoding an MHC antigen of the donor into a bone marrow hematopoietic stem cell of the recipient mammal; and allowing the MHC antigen encoding DNA to be expressed in the recipient.

Preferred embodiments include those in which: the cell is removed from the recipient prior to the DNA insertion and returned to the recipient after the DNA insertion; the DNA includes a MHC class I gene; the DNA includes a MHC class II gene; the DNA is inserted into the cell by transduction, e.g., by a retrovirus, e.g., by a Moloney-based retrovirus; and the DNA is expressed in bone marrow cells and/or peripheral blood cells of the recipient 14, preferably 30, more preferably 60, and most preferably 120 days, after the DNA is introduced into the recipient.

Tolerance, as used herein, refers to the inhibition of a graft recipient's ability to mount an immune response which would otherwise occur, e.g., in response to the introduction of a nonself MHC antigen into the recipient. Tolerance can involve humoral, cellular, or both humoral and cellular responses.

Bone marrow hematopoietic stem cell, as used herein, refers to a bone marrow cell which is capable of developing into mature myeloid and/or lymphoid cells.

- 6 -

MHC antigen, as used herein, refers to a protein product of one or more MHC genes; the term includes fragments or analogs of products of MHC genes which can evoke an immune response in a recipient organism.

- 5 Examples of MHC antigens include the products (and fragments or analogs thereof) of the human MHC genes, i.e., the HLA genes. MHC antigens in swine, e.g., miniature swine, include the products (and fragments and analogs thereof) of the SLA genes, e.g., of the DRB gene.
- 10 Miniature swine, as used herein, refers to partially inbred miniature swine.

Graft, as used herein, refers to a body part, organ, tissue, or cells.

- The invention allows the reconstitution of a graft
- 15 recipient's bone marrow with transgenic autologous bone marrow cells expressing allogeneic or xenogeneic MHC genes. Expression of the transgenic MHC genes confers tolerance to grafts which exhibit the products of these or closely related MHC genes. Thus, methods of the
- 20 invention provide for the induction of specific transplantation tolerance by somatic transfer of MHC genes.

- Methods of the invention avoid the undesirable side effects of broad spectrum immune suppressants which
- 25 are often used in transplantation. Drugs such as Prednisone, Imuran, CyA, and most recently FK506, have all had important impact on the field of transplantation. However, all of these drugs cause nonspecific suppression of the immune system which must be titrated sufficiently
- 30 to avoid rejection while not completely eliminating immune function. Patients must stay on chronic immunosuppressive therapy for the remainder of their lives, facing the major complications of too much or too little immunosuppression, infection and rejection,
- 35 respectively.

- 7 -

Tolerance to transplantation antigens can be achieved through induction of lymphohematopoietic chimerism by bone marrow transplantation (BMT). However, successful BMT across MHC barriers has two major risks:

5 if mature T cells are not removed from the marrow inoculum the recipient may develop severe graft versus host disease (GVHD); removal of these cells often leads to failure of engraftment. Methods of the invention, which induce specific tolerance by reconstitution of the

10 recipients bone marrow with autologous (as opposed to allogeneic or heterologous) bone marrow cells, allow tolerance to be conferred with minimal risk of GVHD and with minimal need to remove T cells from the marrow inoculum.

15 Other features and advantages of the invention will be apparent from the following detailed description and from the claims.

Detailed Description

The drawings are first briefly described.

20 Drawings

Fig. 1 is a diagram of the GS4.5 retroviral construct.

Fig. 2 is a diagram of the GS4.5 proviral genome and the expected transcripts.

25 Fig. 3 is a representation of flow cytometry profile of transduced cells.

Fig. 4 is a diagram of the transduction assay.

Fig. 5 is a diagram of genetic maps of the C57BL/10, B10.AKM, and B10.MBR strains.

30 Fig. 6 is a diagram of the FACS profile of spleen cells from a recipient of transduced bone marrow.

Fig. 7 is a graph of survival versus time in skin graft experiments.

Fig. 8 is a diagram of FACS analysis of thymocytes

35 from graft rejectors, receptors, and controls.

- 8 -

Fig. 9 is a diagram of the N2-B19-H2b vector.

MHC Genes

MHC genes for a variety of species are well studied. For example the HLA genes in man, see, e.g.,
5 Hansen et al., 1989, *The Major Histocompatibility Complex*, In *Fundamental Immunology* 2d ed., W.E. Paul, ed., Raven Press Ltd., New York, hereby incorporated by reference, and the SLA genes in swine, see e.g., Sachs
10 et al., 1988, *Immunogenetics* 28:22-29, hereby incorporated by reference, have been cloned and characterized.

A gene encoding a MHC antigen can be used in methods of the invention to confer tolerance to a graft which displays that or a closely related MHC antigen.
15 Methods of the invention can be used to confer tolerance to allogeneic grafts, e.g., wherein both the graft donor and the recipient are humans, and to xenogeneic grafts, e.g., wherein the graft donor is a nonhuman animal, e.g., a swine, e.g., a miniature swine, and the graft recipient
20 is a human.

The individual which supplies the MHC genes should be as genetically similar as possible, particularly in terms of the MHC genes, to the individual which supplies the graft. For example, in allogeneic grafts wherein the
25 implant donor is a human and the implant recipient is a human it is preferable to use MHC genes from the donor. In this embodiment, MHC probes, e.g., a probe from a third person or a synthetic consensus probe, can be used to isolate DNA encoding the MHC gene or genes of the
30 implant donor individual. This allows the closest match between the genes used to confer tolerance and the genes which express MHC antigens on the graft.

In xenogeneic grafts, the implant donor individual and the individual which supplies the tolerance
35 conferring DNA should be the same individual or should be

- 9 -

as closely related as possible. For example, it is preferable to deriv implant tissue from a colony of donors which is highly inbred and, more preferably, which is homozygous for the MHC genes. This allows the single
5 cloned MHC sequence to be used for many graft recipients.

Transformation of bone marrow cells

MHC genes can be introduced into bone marrow cells by any methods which allows expression of these genes at a level and for a period sufficient to confer tolerance.
10 These methods include e.g., transfection, electroporation, particle gun bombardment, and transduction by viral vectors, e.g., by retroviruses.

Recombinant retroviruses are a preferred delivery system. They have been developed extensively over the
15 past few years as vehicles for gene transfer, see e.g., Eglitis et al., 1988, *Adv. Exp. Med. Biol.* 241:19. The most straightforward retroviral vector construct is one in which the structural genes of the virus are replaced by a single gene which is then transcribed under the
20 control of regulatory elements contained in the viral long terminal repeat (LTR). A variety of single-gene-vector backbones have been used, including the Moloney murine leukemia virus (MoMuLV). Retroviral vectors which permit multiple insertions of different genes such as a
25 gene for a selectable marker and a second gene of interest, under the control of an internal promoter can be derived from this type of backbone, see e.g., Gilboa, 1988, *Adv. Exp. Med. Biol.* 241:29.

The elements of the construction of vectors for
30 the expression of a protein product, e.g., the choice of promoters is known to those skilled in the art. The most efficient expression from retroviral vectors is observed when "strong" promoters are used to control
transcription, such as the SV 40 promoter or LTR
35 promoters, reviewed in Chang et al., 1989, *Int. J. Cell*

- 10 -

Cloning 7:264. These promoters are constitutive and do not generally permit tissue-specific expression.

However, in the case of class I genes, which are normally expressed in all tissues, ubiquitous expression is
5 acceptable for functional purposes. Housekeeping gene promoters, e.g., the thymidine kinase promoter, are appropriate promoters for the expression of class II genes.

The use of efficient packaging cell lines can
10 increase both the efficiency and the spectrum of infectivity of the produced recombinant virions, see Miller, 1990, *Human Gene Therapy* 1:5. Murine retroviral vectors have been useful for transferring genes efficiently into murine embryonic, see e.g., Wagner et al., 1985, *EMBO J.* 4:663; Griedley et al., 1987 *Trends Genet.* 3:162, and hematopoietic stem cells, see e.g., Lemischka et al., 1986, *Cell* 45:917-927; Dick et al., 1986, *Trends in Genetics* 2:165-170.

A recent improvement in retroviral technology
20 which permits attainment of much higher viral titers than were previously possible involves amplification by consecutive transfer between ecotropic and amphotropic packaging cell lines, the so-called "ping-pong" method, see e.g., Kozak et al., 1990, *J. Virol.* 64:3500-3508;
25 Bodine et al., 1989, *Prog. Clin. Biol. Res.* 319:589-600.

Transduction efficiencies can be enhanced by pre-selection of infected marrow prior to introduction into recipients, enriching for those bone marrow cells expressing high levels of the selectable gene, see e.g.,
30 Dick et al., 1985, *Cell* 42:71-79; Keller et al., 1985, *Nature* 318:149-154. In addition, recent techniques for increasing viral titers permit the use of virus-containing supernatants rather than direct incubation with virus-producing cell lines to attain efficient
35 transduction, see e.g., Bodine et al., 1989, *Prog. Clin.*

- 11 -

- Biol. Res. 319:589-600. Because replication of cellular DNA is required for integration of retroviral vectors into the host genome, it may be desirable to increase the frequency at which target stem cells which are actively cycling e.g., by inducing target cells to divide by treatment in vitro with growth factors, see e.g., Lemischka et al., 1986, *Cell* 45:917-927, a combination of IL-3 and IL-6 apparently being the most efficacious, see e.g., Bodine et al., 1989, *Proc. Natl. Acad. Sci.* 86:8897-8901, or to expose the recipient to 5-fluorouracil, see e.g., Mori et al., 1984, *Jpn. J. Clin. Oncol.* 14 Suppl 1:457-463, prior to marrow harvest, see e.g., Lemischka et al., 1986, *Cell* 45:917-927; Chang et al., 1989, *Int. J. Cell Cloning* 7:264-280.
- 15 N2A or other Moloney-based vectors are preferred retroviral vectors for transducing human bone marrow cells.

Preparative Regimen For The Introduction of Transformed Bone Marrow Cells

- 20 To prepare for bone marrow cells the recipient must undergo an ablation of the immune response which might otherwise resist engraftment.
- The preparative regimens necessary to permit engraftment of modified autologous hematopoietic stem cells may be much less toxic than those needed for allogeneic bone marrow transplantation -- preferably requiring only depletion of mature T cells with monoclonal antibodies, as has been recently demonstrated in a mouse model, see Sharabi et al., 1989, *J. Exp. Med.* 169:493-502. It is possible that transient expression may be sufficient to induce tolerance, which may then be maintained by the transplant even if expression on hematopoietic cells is lost, as has been observed for heart transplants in a mixed xenogeneic bone marrow

- 12 -

transplant model, Ildstad et al., 1985, *Transplant. Proc.* 17:535-538.

Example 1: Sustained expression of a swine class II gene in murine bone marrow hematopoietic cells by retroviral-mediated gene transfer

Overview The efficacy of a gene transfer approach to the induction of transplantation tolerance in miniature swine model was shown by using double-copy retroviral vectors engineered to express a drug-resistance marker (neomycin) and a swine class II DRB cDNA. Infectious particles containing these vectors were produced at a titer of $>1 \times 10^6$ G418-resistant colony-forming units/ml using both ecotropic and amphotropic packaging cell lines. Flow cytometric analysis of DRB-transfected murine fibroblasts subsequently transduced with virus-containing supernatants demonstrated that the transferred sequences were sufficient to produce DR surface expression. Cocultivation of murine bone marrow with high-titer producer lines leads to the transduction of 40% of granulocyte/macrophage colony-forming units (CFU-GM) as determined by the frequency of colony formation under G418 selection. After nearly 5 weeks in long-term bone marrow culture, virus-exposed marrow still contained G418-resistant CFU-GM at a frequency of 25%. In addition, virtually all of the transduced and selected colonies contained DRB-specific transcripts. These results show that a significant proportion of very primitive myelopoietic precursor cells can be transduced with the DRB recombinant vector and that vector sequences are expressed in the differentiated progeny of these cells. These experiments are described in detail below.

Construction and Screening of SLA-DRB Recombinant Retroviruses As in man, Lee et al., 1982, *Nature* 299:750-752, Das et al., 1983, *Proc. Natl. Acad. Sci. USA* 80:3543-3547, the sequence of the swine DRA gene is minimally polymorphic. Therefore, transduction of

- 13 -

alogeneic DRB cDNAs into bone marrow cells should be sufficient to allow expression of allogeneic class II DR molecules on cells committed to express this antigen.

Details of retroviral constructs are given in Fig. 1. Two types of retroviral constructs, GS4.4 and GS4.5, were prepared. The diagram in Fig. 1 depicts the GS4.5 retroviral construct. The arrows in Fig. 1 indicate the directions of transcription. In GS4.5, the orientation of DRB cDNA transcription is the same as viral transcription. In GS4.4 (not shown), the TK promoter and the DRB cDNA were inserted into the 3' LTR of N2A in the reverse orientation of transcription with respect to viral transcription and the simian virus 40 3' RNA processing signal was added. pBSt refers to Bluescript vector sequence (Stratagene). The thymidine kinase (TK) promoter was contained within the 215-base-pair (bp) *Pvu II*-*Pst I* fragment from the herpes simplex virus TK gene, McKnight, 1980 *Nucleic Acids Res.* 8:5949-5964. The simian virus 40 3' RNA processing signal was contained within the 142-bp *Hpa I*-*Sma I* fragment from the pBLCAT3 plasmid, Luckow et al., (1987) *Nucleic Acids Res.* 15:5490-5497, (see Fig. 1). Sequence analysis of the junctions of the promoter, the class II cDNA, and the vector sequences confirmed that the elements of the constructs were properly ligated.

These retroviral constructs were transfected into the amphotropic packaging cell line PA317, and transfectants were selected in G418-containing medium. A total of 24 and 36 clones, transfected, respectively, with the GS4.4 and GS4.5 recombinant plasmids, were tested by PEG precipitation of culture supernatants and slot-blot analysis of viral RNA. Of these, 8 and 12 clones were found, respectively, to be positive for DRB, although the DRB signal was consistently weaker for the GS4.4-derived clones. Analysis of genomic and spliced

- 14 -

transcripts from GS4.5 cells by dot-blot analysis of PEG-precipitated particles revealed heterogeneity among viral transcripts in various clones transfected by GS4.5. In one experiment, two clones contained DRB⁺/Neo⁺ viral RNA, two contained DRB⁺/Neo⁻ RNA, two contained DRB⁻/Neo⁺ RNA, and one showed no class II or Neo signal. G418-resistance (G418^r) titer determination of supernatants from DRB-positive clones confirmed that the average titer produced by GS4.5-transfected clones (10^3 - 10^4 CFU/ml) was significantly higher than that of the GS4.4-transfected clones (10^2 - 10^3 CFU/ml). Further transduction experiments were, therefore, conducted with the best clone, named GS4.5 C4, which produced an initial G418^r titer of 3×10^4 CFU/ml.

Plasmid preparation, cloning procedures, DNA sequencing, RNA preparations, Northern blots, and RNA slot blots were performed by standard methods, Sambrook et al., 1989, *Molecular Cloning: A Laboratory Manual* 2nd Ed. (Cold Spring Harbor Lab., Cold Spring Harbor). Final washes of blots were carried out in 0.1 x SSPE (1 x SSPE = 0.18 M NaCl/10 mM sodium phosphate, pH 7.4/1 mM EDTA) at 60°C for 30 min.

The packaging cell lines PA317, Miller et al., 1986, *Mol. Cell. Biol.* 6:2895-2902, GP+E-86, Markowitz et al., 1988, *J. Virol* 62:1120-1124, psiCRIP, Danos et al., 1988, *Proc. Natl. Acad. Sci. USA* 85:6460-6464, and their derivatives were maintained at 37°C in Dulbecco's modified Eagle's medium (DMEM; GIBCO) with 10% (vol/vol) fetal bovine serum (CELlect Silver; Flow Laboratories) supplemented with 0.1 mM nonessential amino acids (Whittaker Bioproducts), antibiotics penicillin (5 units/ml), and streptomycin (5 µg/ml).

Improvement of the Viral Titer of the C4 Clone

Since recent data indicated that supernatants containing high retroviral titers were the best candidates for

- 15 -

transducing bone marrow cells, Bodine et al., 1990, *Proc. Natl. Acad. Sci. USA* 87:3738-3742, the titer of the C4 producer clone was increased by "ping-pong" amplification, Bestwick et al., 1988, *Proc. Natl. Acad. Sci. USA* 85:5404-5408. Supernatant from nearly confluent C4 cultures was used to transduce GP+E-86 ecotropic packaging cells and G418 selection was applied. Forty-eight clones were isolated and screened by PEG precipitation for production of viral particles.

10 Supernatants from 18 of these clones were DRB-positive by dot-blot analysis of viral RNA and had G418^r titers between 0.5 and 3.5×10^4 CFU/ml). One positive clone was then amplified by the ping-pong technique with the amphotropic hygromycin-resistant packaging line psiCRIP.

15 Supernatants from 48 hygromycin-resistant clones were examined for presence of DRB-positive viral RNA by PEG precipitation and their G418^r titers were determined. All of the clones were positive by dot-blot analysis with the DRB probes and produced titers between 1×10^5 and 1×10^7 CFU/ml. Amphotropic clone GS4.5 A4, which produced the highest titer, was tested for the presence of helper virus by the S + L-assay. No replication-competent helper virus was detected.

Amplification of virus titer was achieved by the ping-pong technique. Since there is evidence that psiCRIP packaging cells are less prone to produce helper virus than PA317 when using certain types of vectors, Miller, 1990, *Hum. Gene Therapy* 1:5-14, DRB recombinant virions were prepared using the psiCRIP/GP-E-86 producer combination. Titer values $> 1 \times 10^7$ CFU/ml with no detectable amphotropic helper viruses were obtained, confirming that this strategy produced safe viral particles suitable for *in vivo* experiments.

Northern blot analysis of GS4.5-producing clones C4, A9, and A4, each derived from a different packaging

- 16 -

cell line, showed a conserved hybridization pattern. RNA species corresponding to the full-length viral genome, the spliced Neo transcript, and the DRB transcription unit were observed with additional RNA species. High molecular size species observed in these experiments may constitute a read-through transcript starting from the TK promoter and ending in the other long terminal repeat (LTR). In contrast to many of the virion-producer clones obtained by transfection that presented erratic DRB transcripts, those obtained by transduction showed stable DRB hybridization patterns suggesting that no recombination events occurred during the amplification procedure.

Retroviral titers were determined as follows.

- 15 Replication-defective retroviral particles were produced from packaging cell lines initially transfected with recombinant construct using the standard calcium phosphate precipitation method, Wigler et al., 1978, *Cell* 14:725-733. Retrovirus production was estimated by the drug-resistance titer (G418-resistant colony-forming units/ml, CFU/ml) as described, Bodine et al., 1990, *Proc. Natl. Acad. Sci. USA* 87:3738-3742. Except for the psiCRIP line, G418 (GIBCO) selection was carried out in active component at 500 μ g/ml for 10-12 days. Hygromycin B selection was applied to psiCRIP-derived packaging clones in medium containing active drug at 50 μ g/ml for 10 days. Replication-competent helper virus titer was assayed on PG4 feline cells by the S⁺L⁻ method, Bassen et al., 1971, *Nature* 229:564-566.
- 30 PEG precipitation of viral particles was performed as follows. Virions contained in 1 ml of culture supernatant were precipitated with 0.5 ml of 30% (wt/vol) polyethylene glycol (PEG) for 30 min. at 4°C. After centrifugation, the pellets were treated with a mixture of RNase inhibitors (vanadyl ribonuclease complex, BRL),
- 35

- 17 -

phenol/chloroform-extracted, and ethanol-precipitated. Pellets were then resuspended in 15.7% (vol/vol) formaldehyde and serial dilutions were dotted onto nitrocellulose membrane.

5 Analysis of DRB Transcription in Packaging Cell

Clones To test for accurate transcription of the introduced DRB cDNA within the different producer clones, Northern blots containing RNAs isolated from these clones were hybridized with the DRB and Neo probes. Fig. 2

10 depicts the structure of the provirus genome and the expected sizes of transcripts initiated from either the viral LTR or the TK promoters. Each of the three GS4.5-containing clones, which were derived from PA317 (clone C4), GP + E-86 (clone A9), and psiCRIP (clone A4) cells, 15 showed DRB-positive transcripts. As reported, Hantzopoulos et al., 1989, *Proc. Natl. Acad. Sci. USA* 86:3519-3523, the unspliced genomic RNA (band a) and the spliced Neo transcript (band b) were observed. In addition, a transcript uniquely hybridizable with the DRB 20 probe was detected that corresponds to the size predicted (1700 bases, band c) for the DRB cDNA transcription unit.

Surface Expression of the SLA-DR Antigen on

Transduced Fibroblasts An in vitro assay was developed to examine surface expression of the SLA-DR antigen on 25 murine fibroblasts. Flow cytometry (FCM) profiles shown in Fig. 3 demonstrate that G418^r titers of 3×10^4 (clone C4) were sufficient to promote expression of the DR antigen on the cell surface of transduced DRA

transfectants. In Fig. 3 solid lines indicate DR cell 30 surface expression (anti-DR antibody binding) (22% and 75% of the bulk population of cells 3 days after transduction with GS4.5 C4, (B) and GS4.5 A4 (C), respectively); dashed lines indicate anti-mouse class I antibody binding (positive control); dotted lines 35 indicate anti-pig CD8 antibody binding (negative

- 18 -

control). Twenty-two percent of the bulk population of transduced cells were DR-positive and subclones maintained class II expression for more than 5 months. The increase in titer (clone A4) correlated with an
5 increase in the number of cells transduced (75% of the transduced population was DR-positive) and with the brightness of the DR signal.

The class II transduction assay was performed as diagramed in Fig. 4. NIH 3T3 cells were transfected with
10 the SLA-DRA^d cDNA inserted in a plasmid expression vector, Okayama et al., 1982, *Mol. Cell. Biol.* 2:161-170. Approximately 3×10^4 cells of a stable DRA transfectant (clone 11/12.2F) that expressed a high level of DRA mRNA were then transduced overnight with 1 ml of DRB-
15 containing retroviral supernatant. Cells were subsequently cultivated in fresh DMEM supplemented with 10% fetal bovine serum and antibiotics for 2 additional days and examined for cell surface expression of the DR antigen by FCM analysis.

20 The class II transduction assay described here provides a fast and simple method to test both the expression and functional titer of retroviral constructs. By using cells transfected with DRA, the need for lengthy double selection after transduction by two separated
25 vectors, Yang et al., 1987, *Mol. Cell Biol.* 7:3923-3928; Korman et al., 1987, *Proc. Natl. Acad. Sci. USA* 84:2150-2154, is obviated. Cell-surface expression of DR heterodimers was demonstrated by FCM analysis 3 days after transduction, providing direct evidence that the
30 transferred sequences were sufficient to produce significant level of DR β chain. More importantly, this test allows determination of "functional" titers based on the expression of the gene of interest rather than on that of the independently regulated drug-resistance
35 marker.

- 19 -

The SLA-DRB probe was an *EcoRI* cDNA fragment containing the complete coding sequence of the DR β chain, Gustafsson et al., 1990, *Proc. Natl. Acad. Sci. USA* **87**:9798-9802. The neomycin phosphotransferase gene (Neo) probe was the *Bcl I*-*Xho I* fragment of the N2A retroviral plasmid, Hantzopoulos et al., 1989, *Proc. Natl. Acad. Sci. USA* **86**:3519-3523.

Expression of Porcine DRB cDNA Transduced into Murine Bone Marrow Progenitor Cells The efficiency with which myeloid clonogenic precursors were transduced was determined by assaying for CFU-GM with and without a selecting amount of G418 after exposure of bone marrow cells to GS4.5-derived virions. Comparison of the number of colonies that formed in the presence and absence of the drug, for two experiments, indicated that $\approx 40\%$ of the initial population of myeloid progenitor cells were transduced. The frequency of G418^r CFU-GM was again determined after a sample of the transduced marrow was expanded under long-term culture conditions for 33 days. Twenty-five percent of the progenitors present after 33 days in culture still gave rise to colonies under G418 selection. Colonies of cells arisen from CFU-GM were examined for the presence of DRB-specific transcripts by converting RNA into cDNA and then performing PCR amplification as described herein and in Shafer et al., 1991 *Proc. Natl. Acad. Sci. USA* **88**:9670. A 360-bp DRB-specific product was detected in five of six G418-selected colonies from freshly transduced marrow, whereas all six colonies similarly derived from transduced progenitors present after 33 days in culture were positive. An additional band of 100 bp observed in some of the samples probably reflects the stochastic nature of nonspecific priming events. DRB-specific transcripts were also detected in the bulk population of drug-resistant colonies and in producer cells but were not

- 20 -

detected in controls such as a bulk population of untransduced colonies, fibroblasts used to provide carrier RNA, and a bulk population of transduced colonies processed as above but without reverse transcriptase.

- 5 These latter data demonstrate that the PCR signal was dependent on the synthesis of cDNA, excluding the possibility that provirus, rather than viral message, was responsible for the amplified fragment.

- Recent improvements including modifications of the
- 10 virus design, increase of viral titers, use of growth factors to stimulate precursor cells, and selection of stem cells prior to transduction have been shown to improve long-term expression of transduced genes in the hematopoietic compartment, Bodine et al., 1990, *Proc. Natl. Acad. Sci. USA* 87:3738-3742; Bodine et al., 1989, *Proc. Natl. Acad. Sci. USA* 86:8897-8901; Wilson et al., 1990, *Proc. Natl. Acad. Sci. USA* 87:439-443; Kang et al., 1990, *Proc. Natl. Acad. Sci. USA* 87:9803-9807; Bender et al., 1989, *Mol. Cell. Biol.* 9:1426-1434. The experiments
- 20 herein show the applicability of the retroviral gene-transfer technique in achieving expression of major histocompatibility complex class II genes transferred into hematopoietic cells. To determine the efficiency with which developmentally primitive hematopoietic cells
- 25 were transduced, the frequency of G418^r CFU-GM was assessed after expanding infected marrow cells kept for 33 days in long-term cultures. Expression of the exogenous DRB cDNA was also monitored in cells derived from transduced CFU-GM present either immediately after
- 30 infection or after an extended culture period. Virtually all of the colonies individually tested were positive for DRB-specific transcript, suggesting that the DRB recombinant vector is suitable for expression in murine hematopoietic cells.

- 21 -

Bone marrow cells were obtained from the femora of 6- to 12-week-old female C57BL/10 mice and were prepared as described, Ildstad et al., 1984, *Nature* 307:168-170. Methylcellulose colony assays for granulocyte/macrophage colony-forming units (CFU-GM), Eaves et al., 1978, *Blood* 52:1196-1210, were performed as described using 5% (vol/vol) murine interleukin 3 culture supplement (Collaborative Research). Long-term Dexter-type bone marrow cultures were initiated in 60-mm culture dishes with 2×10^7 nucleated cells, Eaves et al., 1987, *CRC Crit. Rev. Oncol. Hematol.* 7:125-138.

Bone marrow cells were transduced essentially as described, Bodine et al., 1989, *Proc. Natl. Acad. Sci. USA* 86:8897-8901. Briefly, bone marrow was harvested for 6- 12-week-old female C57BL/10 donors that had been treated 2 days with 5-fluorouracil (150 mg/kg). Prestimulation was performed by incubating 1×10^6 cells per ml for 2 days in long-term Dexter-type bone marrow culture medium to which was added 7.5% interleukin 3 culture supplement and recombinant human interleukin 6 (200 units/ml; gift from J. Jule, National Institutes of Health, Bethesda, MD). Marrow cells were transduced for 48 hr by adding 5×10^6 cells per 10-cm plate containing nearly confluent virus-producers, Polybrene (8 mg/ml), and the cytokines described above.

Detection of DRB-Specific Transcripts in CFU-Derived Colonies was performed as follows. Cells corresponding to individual CFU colonies and to colonies present on an entire plate (bulk) were first extracted from methylcellulose cultures by dilution in phosphate-buffered saline and centrifugation. These cells were then combined with 1×10^6 NIH 3T3 cells (to provide carrier RNA), and total RNA was prepared using the guanidine isothiocyanate/CsCl method. First-strand cDNA was prepared from 20 μ g of total RNA using the Invitrogen

- 22 -

Red Module kit. cDNA was then subjected to 50 cycles of PCR amplification in the presence of the SLA DRB-specific oligonucleotides 04 (5'-CCACAGGCCTGATCCCTAATGG) (Seq.

I.D. No. 1) and 17 (5'-AGCATAGCAGGAGCCTTCTCATG) (Seq.

- 5 I.D. No. 2) using the Cetus GeneAmp kit as recommended (Perkin-Elmer/Cetus). Reaction products were visualized after electrophoresis on a 3% NuSieve agarose gel (FMC) by staining with ethidium bromide.

- FCM analysis was performed with a FAC-Scan II
10 fluorescence-activated cell sorter (Becton Dickenson) on cells stained with the anti-DR monoclonal antibody 40D, Pierres et al., 1980, *Eur. J. Immunol.* 10:950-957, an anti-H-2^d allo antiserum, or the anti-porcine CD8 monoclonal antibody 76-2-11, Pescovitz et al., 1984, *J.*
15 *Exp. Med.* 160:1495-1505, followed by fluorescein isothiocyanate-labeled goat anti-mouse antibodies (Boehringer Mannheim).

Example 2: Expression of Allogeneic Class II cDNA in Swine Bone Marrow Cells Transduced With A Recombinant

20 Retrovirus

- A MHC gene (DRB) was transferred into clonogenic progenitor cells from swine using a recombinant retroviral vector (GS4.5) and a transduction protocol designed to be applicable in vivo. Both the selectable
25 drug resistance gene and the allogeneic class II cDNA transferred by this vector were expressed in the progeny of these transduced progenitors. Expression of the Neo gene was monitored functionally by colony formation under G418 selection, while the presence of class II
30 transcripts was detected by PCR analysis. With this latter method, the transcriptional expression of both endogenous and virally derived DRB genes in transduced and selected colonies were demonstrated.

- Primary porcine fibroblasts were cultured with
35 high titer viral supernatants, and then analyzed by

- 23 -

northern blotting using probes specific for DRB and Neo. A specific transcript was observed which was uniquely hybridizable with the DRB probe and migrated at the position predicted (1700 bases) for the DRB cDNA

- 5 transcription unit arising from the TK promoter and terminating at the LTR 3' RNA processing site.

To determine whether GS4.5 containing virions could transduce swine myelopoietic progenitor cells a colony assay adapted for swine CFU-GM was used.

- 10 Transductions were carried out by incubating bone marrow from a donor of the SLA^c haplotype in high titer viral supernatant. Comparisons of the number of colonies which formed in the presence and absence of G418 for a total of 5 independent experiments indicated that 5% to 14% of
15 CFU-GM were transduced.

- Colonies of cells originating from transduced CFU-GM were examined for the presence of DRB-specific transcripts by converting RNA into cDNA, and then performing PCR amplification. Utilizing a polymorphic
20 Sau3AI restriction site absent from the endogenous DRB^c gene, the presence of DRB^d-specific transcripts was unambiguously demonstrated. Gel electrophoresis of the PCR product demonstrated that a 183/177 bp doublet indicative of the vector-derived DRB^d transcript was
25 amplified in samples derived not only from pools of transduced and selected CFU-GM progeny, but also from at least 4 out of 6 individual colonies tested. A 360 bp PCR fragment, indicative of endogenous DRB^c transcripts, was also amplified not only as expected from PBL isolated
30 from an SLA^c donor, but also from both of the pooled colony samples and a number of the individual colony samples.

- Construction of the retrovirus GS4.5, and production of high titer viral supernatants was as
35 described above. Detection of DRB-specific transcripts

- 24 -

in CFU-derived colonies by PCR of cDNA were described above and as follows. Bone marrow from an SLA^c donor was exposed to GS4.5-containing virions, and G418 selected colonies were tested for the presence of DRB^c

5 (endogenous) and DRB^d (vector derived) specific transcripts by PCR of cDNA followed by digestion with Sau3AI and agarose gel electrophoresis. Controls were as follows: template synthesized either in the presence or absence of reverse transcriptase; template derived from
10 cells producing GS4.5-containing virions, from PBL isolated from SLA^c or SLA^d donors, and from untransduced producer cells used as carrier RNA.

Transduction of bone marrow was performed as follows. Swine bone marrow was harvested as previously
15 described (Pennington et al., 1988, *Transplantation* 45:21-26) and transductions were carried out by incubating marrow cells in high titer viral supernatants at an m.o.i. of 3-5 in the presence of 8 ug of polybrene per ml at 37°C for 5 hr. Myeloid progenitors were
20 assayed by colony formation in methylcellulose cultures using PHA-stimulated swine lymphocyte conditioned medium as a source of growth factors. Selective medium contained 1.2 mg/ml active G418.

Transduced bone marrow was administered to a
25 lethally irradiated miniature swine. At 5 weeks peripheral blood lymphocytes were analyzed by Southern, northern, and cell-surface FACS analyses. By all of these test there was evidence of presence of the transduced allogeneic class II gene in these cells and
30 for expression of the product of this gene. In particular, northern analysis showed bands characteristic of the transcribed cDNA, and FACS analysis with a combination of alloantisera and monoclonal antibodies to DR showed presence of the transduced allele of DR beta on
35 the surface of peripheral lymphocytes.

- 25 -

Example 3: Allogenic Tolerance

Development of the B10.MBR - B10.AKM Strain

Combination In an attempt to maintain strains which are truly congenic for the MHC, a program of continuous
5 backcrossing of each congenic line to a common background partner strain was instituted more than 15 years ago. Backcross animals were intercrossed and appropriate progeny selected by serologic typing in order to reestablish each congenic line. Thus, C57BL/10 was used
10 as one reference background strain and all other congenic lines on the B10 background were backcrossed once every six to ten generations to this C57BL/10 line.

During the backcrossing of each congenic line to its pedigreed reference line, there is of course the
15 chance for an intra-MHC recombination event to occur. Typing of the intercross (F_2) generation serologically reveals such recombinant events, and when the recombinant provides a new haplotype of potential interest for genetic studies, it is outcrossed and then intercrossed
20 to produce a homozygous new recombinant H-2 haplotype. One of the most valuable of such recombinants originating in this colony is the B10.MBR line, Sachs et al., 1979, *J. Immunol.* 123:1965-1969, which was derived from a recombination event during the backcrossing of B10.AKM to
25 C57BL/10. Because this strain was the first to separate K^b from I^k it has been used extensively in studies of H-2 immunogenetics. In addition, in combination with the parental B10.AKM strain, the B10.MBR offers the possibility of examining an isolated K gene as the only
30 MHC difference between these two strains. Thus, as illustrated in Fig. 5, introduction of the K^b gene into B10.AKM bone marrow stem cells, could theoretically lead to expression of all cell surface MHC antigens of the B10.MBR. Expression on bone marrow derived cell
35 populations produces transplantation tolerance to the

- 26 -

product of the transduced gene, and this tolerance can be tested by a tissue graft from the B10.MBR strain.

Reconstitution of Myeloablated Mice with Transduced Bone

Marrow Eighty prospective donor B10.AKM mice were

5 treated with 150 mg/kg 5FU on day -7. Bone marrow was harvested from these mice on day -5, treated with anti-CD4 and anti-CD8 monoclonal antibodies (mAbs) plus complement to remove mature T cells, and cultured for five days with N2-B19-H2b virus-containing supernatant
10 (H2) from the psi-Crip packaging cell line. As a control, one-half of the marrow was cultivated with supernatant from control packaging cells not containing N2-B19-H2b (A2). On day zero, 45 B10.AKM recipients received 10 Gy total body irradiation (TBI), followed by
15 administration of various concentrations of cultured bone marrow cells (A2 or H2).

K^b expression On day 13 several animals receiving the lowest doses of cultured bone marrow were sacrificed and individual spleen colonies were harvested and
20 analyzed by PCR for the presence of N2-B19-H2b DNA. In addition, spleen cell suspensions were prepared and analyzed for cell surface expression of K^b by flow microfluorometry on a fluorescence-activated cell sorter (FACS). FACS analyses indicated that all animals
25 receiving the H2-treated marrow showed some level of K^b expression above control staining with the non-reactive antibody. The results are shown in Fig. 6 which is a FACS profile of spleen cells from a recipient of transduced bone marrow: A=Anti K^b antibody; B=control
30 antibody. Spleen cells from recipients of non-transduced marrow were also negative. In addition, the PCR analysis showed every colony examined to contain the transduced DNA. Animals were thereafter followed by FACS and PCR on peripheral blood lymphocytes (PBL). On day 28 and again
35 on day 40, PCR analyses were positive. However, FACS

- 27 -

analysis for cell-surface expression was variable, with PBL from most H2 animals showing only a slight shift of the entire peak for staining with anti-K^b, as compared to PBL from A2 animals stained with the same antibody, or as
5 compared to PBL from H2 animals stained with the non-reactive HOPC antibody.

Allogeneic grafts On day 40 skin from B10.MBR (K^b specific) and B10.BR (control, third party class I disparate) donors was grafted onto all animals. Graft
10 survivals were scored daily by a blinded observer (i.e., readings were made without knowledge of which graft was from which donor strain) until rejection was complete. The survival times are shown in Fig. 7, and indicate marked specific prolongation of survival of the B10.MBR
15 skin grafts on the recipients of K^b-transduced BMC (Fig. 7B), but not on recipients of control marrow (Fig. 7A). One of the animals with a long-standing intact B10.MBR skin graft was sacrificed at day 114 and cell suspensions of its lymphoid tissues were examined by FACS and
20 compared to similar suspensions of cells from an animals which had rejected its B10.MBR skin graft. A striking difference was noted in staining of thymus cells with an anti-K^b mAb. Cell suspensions were prepared and stained either with the anti-K^b mAb 28-8-6 or the control
25 antibody HOPC1. A subpopulation of thymus cells from the tolerant animal showed a marked shift toward increased staining with 28-8-6 compared to HOPC1, while there was essentially no change in the staining pattern of thymocytes from the animal which had lost its graft.
30 Fig. 8 shows FACS analysis on thymocytes from skin graft rejector (Fig. 8A, B) and skin graft acceptor (Fig. 8C, D). Staining with control HOPC1 antibody (Fig. 8A, C) and with specific anti-K^b antibody (Fig. 8B, D). A similar comparison of staining patterns on bone marrow
35 cells showed the presence of low level K^b expression on a

- 28 -

cell population in the marrow of the tolerant mouse, but not of the mouse which had rejected its skin graft. These results indicate that a pluripotent stem cell or early progenitor cell population expressed K^b in the tolerant mouse but not in the rejector mouse, and that this BMC stem cell provided a continuous source of K^b antigen in the thymus on cells which are critical for the inactivation of developing thymocytes with K^b-reactive TCR. It is of interest to note that K^b expression was not detected on splenocytes of the tolerant mouse, and that, in general, splenocyte expression did not correlate with skin graft tolerance. Since the spleen contains T cells which mature in the thymus, these results suggest that either thymocytes lose expression of K^b as they mature, or that the K^b-bearing thymocytes of this animal were cells of a non-lymphoid lineage, such as macrophages.

Long-term expression As discussed above, the B10.AKM and B10.MBR congenic mouse strains are identical except in the MHC class I region. A recombinant retrovirus containing the class I gene from the B10.MBR strain (H-2K^b) linked to a B19 parvovirus promoter (B19-H2K^b) and a neomycin resistance (neo^r) gene was introduced into B10.AKM (H-2K^k) marrow cells. As a control, a recombinant retrovirus containing only the neo^r gene was introduced into B10.AKM marrow cells. The transduced marrow was injected into lethally irradiated AKM recipients pre-treated with an anti-CD8 monoclonal antibody. Twelve weeks post BMT, quantitative PCR was used to show that the B19-H-2K^b proviral sequences were present in 5%-30% of peripheral blood cells in all recipient animals. Reverse transcriptase PCR was used to demonstrate the B19-H-2K^b mRNA in RNA isolated from bone marrow and spleen of a subset of recipient animals.

- 29 -

Construction of the K^b Retroviral Vector The retr viral vectors used the Maloney murine leukemia virus based vector N2, Armentano et al., 1987, *J. Virol.* 61:1647-1650. The coding regions within this virus were
5 deleted during its construction, and replaced with the selectable marker gene, neomycin phosphotransferase (Neo), which is transcribed from the viral LTR promoter, and provides drug resistance to G418. This conventional N2 virus was then further modified by insertion of a
10 parvovirus-derived promoter, B19, Liu et al., 1991, *J. Virol.* (In Press), downstream from Neo, followed by 1.6 kb of cDNA coding for the class I antigen H-2K^b to form the new recombinant virus N2-B19-H2b. Fig. 9 depicts the N2-B19-H26 retroviral vector: P=PstI; X=XhoI; H=HindIII;
15 E=EcoRI; B=BamHI. This latter cDNA was derived by Waneck et al. during the construction of an H-2^b cDNA library for other purposes, Waneck et al., 1987, *J. Exp. Med.* 165:1358-1370.

Viral producer cell lines were developed using the
20 packaging cell lines for amphotropic (psi-Crip), Danos et al., 1988, *Proc. Natl. Acad. Sci.* 85:6460-6464, and ecotropic (psi-2), Sambrook et al., 1989, *Molecular cloning: A laboratory manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, viral production.
25 These cell lines have been specially designed to produce structural viral proteins for the recombinant defective virus to be produced. Viral production was achieved by transfecting psi-Crip with N2-B19-H2b. Both amphotropic and ecotropic producer cell lines were then co-cultivated
30 allowing multiple integration events and high expression [i.e. the "ping-pong" technique see Bestwick et al., 1988, *Proc. Natl. Acad. Sci.* 85:5404-5408]. In this technique, co-cultivation overcomes viral antigen receptor blockage by endogenously secreted proteins since
35 amphotropic and ecotropic viruses recognize different

- 30 -

receptors. Ecotropic psi-2 viral producer clones were then selected which produced titers of G418 resistance in 3T3 cells of greater than 10^7 cfu/ml.

In order to ensure that K^b was being expressed from the recombinant virus, transduced 3T3 cells were stained with a monoclonal antibody specific to this antigen and analyzed by flow microfluorometry. These experiments clearly demonstrated high level expression of virally derived K^b .

10 Animals and husbandry were as follows. The B10.BMR strain, [Sachs et al., 1979, *J. Immunol.* 123:1965-1969, was provided to the Jackson Laboratory, Bar Harbor, ME about 6 years ago, and specific pathogen-free stock animals of this strain are now available from that source. Upon arrival in animals should be transferred to autoclaved microisolator cages containing autoclaved feed and autoclaved acidified drinking water. Sterile animal handling procedures which are effective in maintaining animals free of pathogens so that interpretable survival studies can be performed should be used.

20 Bone marrow transplantation was performed as follows. Techniques for bone marrow transplantation in mice are known to those skilled in the art, see e.g., Sykes et al., 1988, *J. Immunol.* 140:2903-2911. Briefly, recipient B10.AKM mice aged 12 to 16 weeks are lethally irradiated (1025R, 137Cs source, 110R/min) and reconstituted within 8 hours with 2.5×10^6 bone marrow cells, obtained from the tibiae and femora of sex-matched donors aged 6-14 weeks. Animals are housed in sterilized microisolator cages, and receive autoclaved food and autoclaved acidified drinking water. For these studies some modifications of this general technique are required, since the syngeneic bone marrow will have been transduced with an allogeneic gene, and since the bone marrow will come from 5FU-treated mice, which should have

- 31 -

lower total cell counts but higher stem cell content than normal mice. The protocol is therefore as follows:

1. Donors will be treated with 5-Fluorouracil, 150 mg/kg i.v. on day -7 in order to induce pluripotent stem cells to cycle.

2. Marrow will be harvested from donors on day -5, and T cell depleted with mAbs and complement.

3. Marrow will then be cultured for 5 days in supernatant from an ecotropic packaging cell line (B17H2Kb-18) which produces a high titer of non-infectious retroviral particles containing the K^b gene (see below). IL-3 and IL-6 will be added to the cultures.

4. On day 0, recipient B10.AKM mice will be lethally irradiated (10.25 Gy), and will be reconstituted with 2.5×10^6 BMC transduced with the K^b gene. Control animals will be similarly treated, except that they will receive marrow exposed to supernatant from a similar ecotropic packaging line not exposed to a K^b -containing vector. The recipient may also be pre-treated with anti-CD8 monoclonal antibody.

Cellular and serological assays are performed as follows.

Anti-class I Cell-Mediated Lympholysis (CML)

25 Assay: Spleens are removed from BMT recipients and normal mice, red cells are lysed using ACK buffer, and a single cell suspension is prepared. Cells are filtered through 100-mesh nylon, washed, and resuspended at 4×10^6 /ml in complete medium consisting of RPMI 1640 with 10% fetal calf serum, 0.025mM 2-mercaptoethanol, .01M Hepes buffer, .09mM nonessential amino acids, 1mM sodium pyruvate, 2mM glutamine, 100U/ml penicillin and 100ug/ml streptomycin. 90 μ l of responder cells are added to Costar 96-well round-bottomed plates along with irradiated (30 Gy) stimulator splenocytes. Cultures are

- 32 -

set up in two rows of 3 replicates each, and after 5 days of incubation in 6% CO₂ at 37°C, twofold serial dilutions are prepared from the second row of triplicates, so that cytolytic capacity can be examined at a total of 5
5 different responder:target ratios. ⁵¹Cr-labelled 2-day concanavalin A-induced lymphoblasts are then added at 10⁴ blasts per well and incubated for 4 hr at 37°C, 6% CO₂. Plates are harvested using the Titertek supernatant collection system (Skatron, Inc., Sterling, VA) and ⁵¹Cr
10 release is determined using an automated gamma counter. Cytolytic capacity is measured directly in the original cell culture plated, so that the measurement is based on the number of responders plated, rather than on the number of live cells present at the end of the 5-day
15 incubation period. This methodology has been developed and used successfully in this laboratory for several years for analysis of spleen cell responses from individual animals [Sykes, M., et al., 1988 J.Immunol. 140:2903-2911]. Percent specific lysis is calculated
20 using the formula:

% Specific Lysis =

$$\frac{\text{Experimental release} - \text{Spontaneous release}}{\text{Maximum release} - \text{Spontaneous release}} \times 100\%$$

Limiting dilution analyses: Responder and
25 stimulator (6x10⁵, 30 Gy irradiated) cells are concultured for 7 days in complete medium containing 13% TCGF (lectin-inactivated con A supernatant obtained from BALB/c con A-activated splenocytes) in 96-well plates. Wells containing 10⁵ (24 wells), 3x10⁴ (24 wells), 10⁴ (30
30 wells), 3000 (30 wells), 1000 (30 wells), 300 (30 wells), and 100 (30 wells) responder cells are prepared. Three thousand ⁵¹Cr-labelled con A blasts are added to each well on day 7, and 4 hour ⁵¹Cr release is measured. Wells are considered positive if ⁵¹Cr release is 3 standard

- 33 -

deviations greater than the mean ^{51}Cr release in 24 wells containing stimulator cells only plus similar numbers of target cells. The Poisson distribution is used to determine the frequency of precursor CTL's which
5 recognize each target, and statistical analysis is performed by the Chi square method of Taswell, Taswell, 1981, *J.Immunol.* 126:1614.

Flow microfluorometry: One-color and two-color flow cytometry will be performed, and percentages of
10 cells expressing a particular phenotype will be determined from 2-color data, as previously described in detail Sykes, 1990, *J.Immunol.* 145:3209-3215. The Lysis II software program (Becton Dickinson) will be used for distinguishing granulocytes from lymphocytes by gating on
15 the basis of forward angle and 90° light scatter. Cell sorting will be performed on a Coulter Epics Elite cell sorter.

Cell suspensions for flow cytometry: PBL, BMC, thymocyte, splenocyte, and lymph node suspensions will be
20 prepared as previously described, Sykes, M. et al., 1988, *J.Immunol.* 140:2903-2911; Sykes, M. 1990, *J.Immunol.* 145:3209-3215; Sharabi, Y. et al., 1990, *J.Exp.Med.* 172:195-202 Whole peripheral white blood cell suspensions (including granulocytes) will be prepared by
25 centrifugation of heparinized blood for 2 minutes at 14,000 RPM in an Eppendorf centrifuge, followed by aspiration of the buffy coat layer. These cells will be transferred to a 15 ml. conical tube and washed. Red blood cells (RBC) contaminating the remaining pellet will
30 be lysed by exposure for 5 seconds to 4.5 ml of distilled H_2O followed by rescue with 0.5 ml of 10x PBS.

Cell staining: One-color and two-color staining will be performed as we have previously described, Sykes, M., 1990, *J. Immunol.* 145:3209-3215; Sykes et al., 1988,
35 *J. Immunol.* 141:2282-2288. Culture supernatant of rat

- 34 -

- anti-mouse RcrR mAB 2.4G2, Unkeless, J.C., 1979, *J. Exp. Med.* 150:580-596, will be used for blocking of non-specific staining due to FcrR binding, whenever only direct staining is used. The following mABs are used:
- 5 biotinylated murine K^b-specific IgG_{2a} mAB 28-8-6, Ozato et al., 1981, *J. Immunol.* 126:317-321, and control murine IgG_{2a} mAB HOPC1 (with no known specific binding to murine antigens) are prepared by purification on a protein A-Sepharose column, and are biotinylated by standard
- 10 procedures used in our laboratory; rat anti-MAC1 mAB M1/70, Springer et al, 1979, *Eur. J. Immunol.* 9:301, is used as culture supernatant, and will be stained by mouse anti-rat IgG-specific mAB MAR18.5; FITC-labelled rat-anti-mouse granulocyte antibody Gr1 is purchased from
- 15 Pharmingen; FITC-labelled rat-anti-mouse IgM mAb is purchased from Zymed; FITC-labelled rat-anti-mouse Thy1.2 mAb will be purchased from Becton-Dickinson; FITC-labelled mouse-anti-human CD3 mAb Leu4 (Becton Dickinson) is used as a directly FITC labelled negative control
- 20 antibody.

- Thymic tissue immunofluorescence: The tissue is incubated in L15 medium for 24 hours to reduce background staining, and is then cut and embedded in O.C.T. compound for freezing in Isopentane. Frozen sections are prepared
- 25 (thickness 4 μ m) on a cryostat, dried, fixed in acetone, then washed in PBS. The first antibody incubation (with 28-8-6) is performed in the presence of 2% normal mouse serum, in order to saturate Fc receptors. After 45 minutes, the slides are washed 4 times, and FITC-
- 30 conjugated secondary reagent (monoclonal rat-anti-mouse IgG_{2a}-FITC, purchased from Pandex) is added. After 45 minutes' incubation with the secondary reagent, four washes are performed and the tissue is mounted. Sections are examined under a fluorescence microscope by an

- 35 -

observer who is unaware of the group of animals from which the tissue was obtained.

Bone Marrow Manipulations and Assays were performed as follows:

- 5 Transduction of murine bone marrow stem cells: The methodology used for transduction of bone marrow cells has been described previously, Karlsson et al., 1988, *Proc. Natl. Acad. Sci.* 85:6062-6066. Bone marrow is harvested from 6-12 week old female B10.AKM donors
- 10 treated 2 days previously with 150 mg/kg 5-FU. Following T cell depletion (see above), the marrow is divided and 10^7 cells per 10 cm plate are cultured for 5 days in the presence of 8 μ g of Polybrene per ml, 10% FCS, 0.6% IL-3-containing supernatant, 0.6% IL-6-containing supernatant,
- 15 and fresh supernatants from B19H2K^b or N2 cells. IL-3- and IL-6-containing supernatant is 48 hour supernatant of COS 7 cells transfected with the murine rIL-3 gene-containing plasmid pCD-IL-3 or with the murine rIL-6 gene-containing plasmid pCD-IL-6, respectively (both
- 20 plasmids provided by Dr. Frank Lee, DNAX Corp.). IL-3-containing supernatants are tittered by testing proliferation of the IL-3-dependent cell line 32D in the presence of dilutions of these supernatants, and IL-6 is tittered in a similar manner using the IL-6-dependent
- 25 line T1165 as the indicator cell line. We will also test the effect of murine SCF on bone marrow transduction, as recently described, Zsebo et al., 1990, *Cell* 63:125-201.

- The virus-containing supernatants are refreshed on a daily basis by harvesting the non-adherent layer of
- 30 each plate, pelleting the cells, and resuspending in freshly harvested filtered virus-containing B19H2K^b or N2 supernatant with additives. After 5 days, the non-adherent and adherent BMC are harvested, washed, and resuspended at 2.5×10^6 /ml in Medium 199 with Hepes buffer

- 36 -

and Gentamycin plus Heparin 10 U/ml. One ml. of this suspension is injected i.v. to irradiated mice.

Murine CFU-GM assay: To test for the bone marrow progenitor cells known as CFU-GM (colony forming unit-
5 granulocyte/macrophage), bone marrow cells are suspended in plating medium consisting of IMDM medium containing 30% defined fetal bovine serum (FBS) (HyClone, Logan, UT), 10^{-4} M β -mercaptoethanol, antibiotics, 5% v/v murine IL-3 culture supplement (Collaborative Research Inc.,
10 Bedford, MA) and 0.8% methylcellulose (achieved by adding 36% v/v of a commercially prepared solution purchased from the Terry Fox Laboratory, Vancouver). 1.1 ml of this suspension is then dispensed into 35mm tissue culture plates (in duplicate), and placed in a 37°C
15 incubator. The resulting CFU-GM derived colonies are enumerated microscopically after 5-7 days. Transduced CFU-GM are selected by including 0.9 ug/ml active G418 in the culture medium. The transduction frequency is then determined by the ratio of CFU-GM which form colonies in
20 the presence and in the absence of the drug.

Molecular methods were as follows:

Construction of N2-B19-H2b vector: This vector was constructed starting from the original retroviral vector N2, Eglitis et al., 1985, *Science* 230:1395-1398, as
25 modified by Shimada to include an additional BamHI site immediately 3' of the XhoI site. It includes the K^b cDNA previously cloned in the vector pBG367, as described by G. Waneck, Waneck et al., 1987, *J. Exp. Med.* 165:1358-1370. This gene has been placed under control of the B19
30 promoter, a highly efficient parvo virus derived promoter, Liu et al., 1991, *J. Virol. In Press*:] to produce the N2-B19-H2b construct.

Southern blot analysis can be performed on DNA extracted from PBL, thymocyte, BMC, splenocyte or lymph
35 node cell suspensions using standard methods, Ausubel et

- 37 -

al., 1989, *Current protocols in molecular biology*. John Wiley & Sons, New York, and probing will be performed with the fragment of K^b cDNA released from pBG367 by EcoRI. The genomic DNA will be cut with enzymes capable
5 of distinguishing the transduced K^b from other class I genes of the B10.AKM strain. From known sequences it would appear that EcoRI may be satisfactory for this purpose, since it should liberate a 1.6 kb band from the transduced K^b cDNA, which is distinct from both the
10 expected endogenous K^k and D^q class I bands of B10.AKM, Arnold et al., 1984, *Nucl. Acids Res.* 12:9473-9485; Lee et al., *J. Exp. Med.* 168:1719-1739. However, to assure that there is no confusion with bands liberated from
15 other class I and class I-like genes we will test several enzymes first on DNA from B10.AKM and choose appropriate restriction enzyme combinations.

PCR analysis of DNA can be performed using primers previously shown to be effective in our preliminary studies (see Fig. 4):

20 5' primer: 5'-GGCCACACTCGCTGAGGTATTTTCGTC-3' (covers 5' end of $\alpha 1$ exon) (Seq. ID No. 3)
3' primer: 5'-GCCAGAGATCACCTGAATAGTGTGA-3' (covers 5' end of $\alpha 2$ exon) (Seq. ID No. 4)

DNA is subjected to 25 cycles of PCR amplification
25 using these specific oligonucleotides and the Cetus GeneAmp kit (Perkin Elmer Cetus, Norwalk, CT) according to the manufacturer's directions. In addition, [³²]PdCTP will be included in the PCR reaction in order to visualize products by autoradiography following
30 electrophoresis.

RNA can be isolated from 5x10⁶ to 5x10⁷ cells using the guanidine isothiocyanate and CsCl methods, Chirgwin et al., 1979, *Biochem.* 18:5294-5308, and will be used for northern analyses, RNase protection analyses,
35 and for PCR analyses of products formed by reverse

- 38 -

transcriptase. For situations in which less than 5×10^6 cells are available, for example following tail bleedings of individual mice, we will utilize the QuickPrep mRNA Purification Kit (Amgen) as a miniaturized RNA

5 preparation procedure.

Northern analyses can be carried out using standard methods, Ausubel et al., 1989, *Current protocols in molecular biology* John Wiley & Sons, New York, and the same K^b cDNA-derived probe. Vector-derived K^b mRNA is
10 larger than endogenous class I transcripts (2.5 kb vs. 1.6 kb) due to the inclusion of vector sequences between the 3' end of the cDNA and the poly-adenylation site in the viral 3' LTR. It should therefore be easy to distinguish the vector-derive K^b mRNA from endogenous
15 transcripts that might cross-hybridize with a K^b cDNA probe. We will also utilize probes derived from unique non- K^b sequences of the transcript (e.g., from B19 or N2 derived vector sequences).

RNAse protection analyses are more sensitive than
20 standard northern blots, yet still quantitative. Procedures based on published methods, Sambrook et al., 1989, *Molecular cloning: A laboratory manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, will be used to derive riboprobes. Briefly, the K^b cDNA will
25 be cloned into a plasmid vector containing the T3 and T7 RNA polymerase promoter sequences (bluescript or Bluescribe plasmids from Stratagene). Using appropriate polymerase and ^{32}P -nucleotides, transcription of the insert will be initiated and the radioactive K^b RNA will
30 be purified. This probe will then be incubated with various RNA preparations followed by treatment with ribonuclease. Presence of RNA will be assessed by electrophoresis on a sequencing gel.

PCR following reverse transcriptase treatment of
35 RNA will be used as a highly sensitive procedure for

- 39 -

detecting the K^b transcript. Appropriate primers will be designed in order to specifically amplify retroviral derived transcripts (one primer covering the 5'UT region of the construct and a second derived from the cDNA
5 sequence). Briefly, RNA will be prepared by the GuSCN/CsCl method and first strand cDNA will be prepared from 5ug of total RNA using the SuperScript preamplification system (BRL/Life Technologies, Inc., Gaithersburg, MD). PCR amplifications will be conducted
10 for 50 cycles, Hansen et al., *J. Immunol.* 118:1403-1408, using the Cetus GeneAmp kit (Perkin Elmer Cetus, Norwalk, CT). Reaction products will be visualized following electrophoresis on a 3% NuSieve agarose gel (FMC BioProducts, Rockland, ME).

- 40 -

SEQUENCE LISTING**(1) GENERAL INFORMATION:**

- (i) APPLICANT: Sachs, David J.
- (ii) TITLE OF INVENTION: SPECIFIC TOLERANCE IN TRANSPLANTATION
- (iii) NUMBER OF SEQUENCES: 4
- (iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Fish & Richardson
(B) STREET: 225 Franklin Street
(C) CITY: Boston
(D) STATE: Massachusetts
(E) COUNTRY: U.S.A.
(F) ZIP: 02110-2804

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: 3.5" Diskette, 1.44 Mb
(B) COMPUTER: IBM PS/2 Model 50Z or 55SX
(C) OPERATING SYSTEM: IBM P.C. DOS (Version 3.30)
(D) SOFTWARE: WordPerfect (Version 5.0)

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:
(B) FILING DATE:
(C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: 07/797,555
(B) FILING DATE: 22 November 1991

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Clark, Paul T.
(B) REGISTRATION NUMBER: 30,162
(C) REFERENCE/DOCKET NUMBER: 00786/111001

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (617) 542-5070
(B) TELEFAX: (617) 542-8906
(C) TELEX: 200154

- 41 -

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

CCACAGGCCT GATCCCTAAT GG 22

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 23
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

AGCATAGCAG GAGCCTTCTC ATG 23

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 3:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 27
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

GGCCCACACT CGCTGAGGTA TTTCGTC 27

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 25
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- 42 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

GCCAGAGATC ACCTGAATAG TGTGA

25

Other embodiments are within the following claims.
What is claimed is:

- 43 -

Claims

1. Use of nucleic acid encoding a mammalian MHC antigen inserted into a bone marrow hematopoietic cell in the preparation of a medicament for conferring tolerance
5 to a transplanted tissue in a recipient mammal.
2. The use of claim 1, wherein said recipient mammal is from a first species and said transplanted tissue is from a second species.
3. The use of claim 2, wherein said mammalian MHC
10 antigen is from said second species.
4. The use of claim 1, wherein said recipient is a human.
5. The use of claim 1, wherein said transplanted tissue is from a swine.
- 15 6. The use of claim 5, wherein said swine is a miniature swine.
7. The use of claim 1, wherein said nucleic acid and said tissue are obtained from the same individual mammal.
- 20 8. The use of claim 1, wherein said nucleic acid is obtained from an individual mammal which is syngeneic to the individual mammal from which said tissue is obtained.
9. The use of claim 1, wherein said nucleic acid
25 is obtained from an individual mammal which is MHC

- 44 -

identical to the individual mammal from which said tissue is obtained.

10. The use of claim 1, wherein said nucleic acid comprises an MHC class I gene.

5 11. The use of claim 1, wherein said nucleic acid comprises an MHC class II gene.

12. The use of claim 1, wherein said nucleic acid is inserted into said cell by transduction.

10 13. The use of claim 12, wherein said nucleic acid is inserted into said cell by a retrovirus.

14. The use of claim 13, wherein said nucleic acid is recipient is a human and said retrovirus is a Moloney-based retrovirus.

15 15. The use of claim 1, wherein said hematopoietic cell is taken from said recipient.

16. The use of claim 1, wherein said recipient mammal is from a first species and said transplanted tissue is from said first species.

20 17. The use of claim 16, wherein said recipient is a human.

18. The use of claim 16, wherein said nucleic acid and said tissue are obtained from the same mammal.

19. The use of claim 16, wherein said nucleic acid is obtained from an individual mammal which is

- 45 -

syngeneic to the individual mammal from which said tissue is obtained.

20. The use of claim 16, wherein said nucleic acid is obtained from an individual mammal which is MHC
5 identical to the individual mammal from which said tissue is obtained.

21. The use of claim 16, wherein said nucleic acid comprises an MHC class I gene.

22. The use of claim 16, wherein said nucleic
10 acid comprises an MHC class II gene.

23. The use of claim 16, wherein said nucleic acid is inserted into said cell by transduction.

24. The use of claim 23, wherein said nucleic acid is inserted into said cell by a retrovirus.

15 25. The use of claim 24, wherein said nucleic acid is recipient is a human and said retrovirus is a Moloney-based retrovirus.

26. The use of claim 16, wherein said hematopoietic cell is taken from said recipient.

20 27. Use of a bone marrow hematopoietic cell comprising nucleic acid encoding a mammalian MHC antigen in the preparation of a medicament for conferring tolerance to a transplanted tissue in a recipient mammal.

25 28. The use of claim 27, wherein said recipient mammal is from a first species and said transplanted tissue is from a second species.

- 46 -

29. The use of claim 27, wherein said mammalian MHC antigen is from said second species.

30. The use of claim 27, wherein said recipient is a human.

5 31. The use of claim 27, wherein said transplanted tissue is from a swine.

32. The use of claim 31, wherein said swine is a miniature swine.

10 33. The use of claim 27, wherein said nucleic acid and said tissue are obtained from the same individual mammal.

15 34. The use of claim 27, wherein said nucleic acid is obtained from an individual mammal which is syngeneic to the individual mammal from which said tissue is obtained.

35. The use of claim 27, wherein said nucleic acid is obtained from an individual mammal which is MHC identical to the individual mammal from which said tissue is obtained.

20 36. The use of claim 27, wherein said nucleic acid comprises an MHC class I gene.

37. The use of claim 27, wherein said nucleic acid comprises an MHC class II gene.

25 38. The use of claim 27, wherein said nucleic acid is inserted into said cell by transduction.

- 47 -

39. The use of claim 38, wherein said nucleic acid is inserted into said cell by a retrovirus.

40. The use of claim 34, wherein said nucleic acid is recipient is a human and said retrovirus is a
5 Moloney-based retrovirus.

41. The use of claim 27, wherein said hematopoietic cell is taken from said recipient.

42. The use of claim 27, wherein said recipient mammal is from a first species and said transplanted
10 tissue is from said first species.

43. The use of claim 42, wherein said recipient is a human.

44. The use of claim 42, wherein said nucleic acid and said tissue are obtained from the same mammal.

15 45. The use of claim 42, wherein said nucleic acid is obtained from an individual mammal which is syngeneic to the individual mammal from which said tissue is obtained.

46. The use of claim 42, wherein said nucleic
20 acid is obtained from an individual mammal which is MHC identical to the individual mammal from which said tissue is obtained.

47. The use of claim 42, wherein said nucleic acid comprises an MHC class I gene.

25 48. The use of claim 42, wherein said nucleic acid comprises an MHC class II gene.

- 48 -

49. The use of claim 42, wherein said nucleic acid is inserted into said cell by transduction.

50. The use of claim 49, wherein said nucleic acid is inserted into said cell by a retrovirus.

5 51. The use of claim 50, wherein said nucleic acid is recipient is a human and said retrovirus is a Moloney-based retrovirus.

52. The use of claim 27, wherein said hematopoietic cell is taken from said recipient.

1/4




FIG 1




FIG 2




FIG. 3a




FIG. 3b




FIG 4

3/4




FIG 5




FIG 6




FIG 7a




FIG 7b

4/4




FIG 8A




FIG 8B




FIG 8C




FIG 8D




FIG 9

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US92/09849

A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) :A61K 48/00

US CL :514/44

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : NONE514/44

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
NONE

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, DIALOG CLASS II MMC, GENE THERAPY

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	NATURE, VOLUME 172, ISSUED 03 OCTOBER 1953, BILLINGHAM <u>ET AL.</u> "ACTIVELY ACQUIRED TOLERANCE OF FOREIGN CELLS", PAGES 603-606, SEE ENTIRE ARTICLE.	1-52
Y	TRENDS IN GENETICS, VOLUME 2, ISSUED JUNE 1986, DICK <u>ET AL.</u> "GENETIC MANIPULATED OF HEMATOPOIETIC STEM CELLS WITH RETROVIRUS VECTORS", PAGES 165-170, SEE ENTIRE ARTICLE.	1-52
Y	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES, VOLUMN 87, ISSUED DECEMBER 1990, GUSTAFSSON <u>ET AL.</u> "STRUCTURE OF MINIATURE SWINE CLASS II DRB GENES: CONSERVATIVE OF HYPERVARIABLE AMINO ACID RESIDUES BETWEEN DISTANTLY RELATED MAMMALIAN SPECIES", PAGES 9798-9802, SEE ENTIRE ARTICLE.	1-52
Y	TRANSPLANTATION, VOLUME 45, NO.1, ISSUED JANUARY 1988, PENNINGTON <u>ET AL.</u> "BONE MARROW TRANSPLANTATION IN MINIATURE SWINE", PAGES 21-26.	1-52

☐

Further documents are listed in the continuation of Box C.

☐

See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be part of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E / earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

07 JANUARY 1993

Date of mailing of the international search report

27 JAN 1993

Name and mailing address of the ISA/
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. NOT APPLICABLE

Authorized officer

SUZANNE ZISKA

Telephone No. (703) 308-0196